

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
25 July 2002 (25.07.2002)

PCT

(10) International Publication Number  
**WO 02/057783 A2**

(51) International Patent Classification<sup>7</sup>: **G01N 33/566**,  
C12Q 1/68, G01N 33/50

(21) International Application Number: PCT/US01/48985

(22) International Filing Date:  
18 December 2001 (18.12.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
0031527.5 22 December 2000 (22.12.2000) GB

(71) Applicants (*for all designated States except US*): **GLAXO GROUP LIMITED** [GB/GB]; Glaxo Wellcome House, Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB). **SMITHKLINE BEECHAM P.L.C.** [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **ANDREWS, John, Leon** [US/US]; GlaxoSmithKline, Five Moore Drive, PO Box 13398, Research Triangle Park, NC 27709 (US). **BRISCOE, Celia, Patricia** [GB/US]; GlaxoSmithKline, Five Moore Drive, PO Box 13398, Research Triangle Park, NC 27709 (US). **IGNAR, Diane, Michelle** [US/US]; GlaxoSmithKline, Five Moore Drive, PO Box 13398, Research Triangle Park, NC 27709 (US). **MUIR, Allison, Isobel** [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB). **SAULS, Howard, Ray Jr.** [US/US];

GlaxoSmithKline, Five Moore Drive, PO Box 13398, Research Triangle Park, NC 27709 (US). **TADAYYON, Mohammad** [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB).

(74) Agents: **LEVY, David, J.** et al.; GlaxoSmithKline, Five Moore Drive, PO Box 13398, Research Triangle Park, NC 27709 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— *without international search report and to be republished upon receipt of that report*

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: METHOD OF SCREENING FOR GPR40 LIGANDS

(57) Abstract: Methods of screening for ligands of the GPR40 receptor are provided, including methods utilizing fatty acid GPR40 ligands.



**WO 02/057783 A2**

## METHOD OF SCREENING FOR GPR40 LIGANDS

### Technical Field

This invention relates to the identification of a ligand for a G-protein coupled receptor (GPCR) and its use in screening methods, and rational drug design, to identify antagonists and agonists of the receptor, to the antagonists and agonists so identified and  
5 their use in therapy.

### Background

The membrane protein gene superfamily of G-protein coupled receptors (GPCRs) has been characterized as having seven putative transmembrane domains. The domains are believed to represent transmembrane  $\alpha$ -helices connected by extracellular and  
10 cytoplasmic loops. G-protein coupled receptors include a wide range of biologically active receptors, such as hormone, viral, growth factor and neuroreceptors.

G-protein coupled receptors have been characterized as including these seven conserved hydrophobic stretches of about 20 to 30 amino acids, connecting at least eight divergent hydrophilic loops. The G-protein family of coupled receptors includes  
15 dopamine receptors which bind to neuroleptic drugs used for treating psychotic and neurological disorders. Other examples of members of this family include, but are not limited to, calcitonin, adrenergic, endothelin, cAMP, adenosine, muscarinic, acetylcholine, serotonin, histamine, thrombin, kinin, follicle stimulating hormone, opsins, endothelial differentiation gene-1 receptor, rhodopsins, odorant, cytomegalovirus  
20 receptors.

Most G-protein coupled receptors have single conserved cysteine residues in each

of the first two extracellular loops which form disulfide bonds that are believed to stabilize functional protein structure. The 7 transmembrane regions are designated as TM1, TM2, TM3, TM4, TM5, TM6, and TM7. TM3 has been implicated in signal transduction.

5           Phosphorylation and lipidation (palmitoylation or farnesylation) of cysteine residues can influence signal transduction of some G-protein coupled receptors. Most G-protein coupled receptors contain potential phosphorylation sites within the third cytoplasmic loop and/or the carboxy terminus. For several G-protein coupled receptors, such as the b-adrenoreceptor, phosphorylation by protein kinase A and/or specific  
10       receptor kinases mediates receptor desensitization.

          For some receptors, the ligand binding sites of G-protein coupled receptors are believed to comprise a hydrophilic socket formed by several G-protein coupled receptor transmembrane domains, which socket is surrounded by hydrophobic residues of the G-protein coupled receptors. The hydrophilic side of each G-protein coupled receptor  
15       transmembrane helix is postulated to face inward and form a polar ligand binding site. TM3 has been implicated in several G-protein coupled receptors as having a ligand binding site, such as the TM3 aspartate residue. TM5 serines, a TM6 asparagine and TM6 or TM7 phenylalanines or tyrosines are also implicated in ligand binding.

          G-protein coupled receptors can be intracellularly coupled by heterotrimeric G-  
20       proteins to various intracellular enzymes, ion channels and transporters. See Johnson, *et al.*, *Endoc. Rev.*, (1989) 10: 317-331). Different G-protein  $\alpha$ -subunits preferentially stimulate particular effectors to modulate various biological functions in a cell. Phosphorylation of cytoplasmic residues of G-protein coupled receptors have been identified as an important mechanism for the regulation of G-protein coupling of some G-  
25       protein coupled receptors. G-protein coupled receptors are found at numerous sites within a mammalian host.

          Over the past 15 years, nearly 150 therapeutic agents targeting 7 transmembrane (7 TM) receptors have been successfully introduced onto the market, thereby establishing their value as therapeutic targets.

### Summary

A first aspect of the present invention is a method of screening a test compound to determine whether the compound is a GPR40 receptor ligand. The method comprises detecting whether the test compound competitively inhibits the binding of a fatty acid  
5 GPR40 ligand to a GPR40 receptor.

A further aspect of the present invention is a method of screening a compound for GPR40 antagonist activity. The method comprises providing a test cell expressing on its surface a GPR40 receptor, and containing a reporter construct comprising a Gq responsive transcriptional element and a reporter gene, where expression of the reporter gene  
10 produces a detectable signal. The test cell surface is exposed to a GPR40 fatty acid agonist under conditions that permit binding of a GPR40 fatty acid agonist, and then exposed to a test compound. Any detectable signal produced by the reporter gene is measured, and compared to the detectable signal that would be expected to be produced if the cell were exposed only to the GPR40 agonist. A decrease in the detectable signal  
15 indicates the test compound is a GPR40 antagonist.

A further aspect of the present invention is a method of screening a compound for GPR40 agonist activity. The method comprises providing a cell expressing on its surface a GPR40 receptor, and containing a reporter construct comprising a responsive transcriptional element and a reporter gene, where expression of the reporter gene  
20 produces a detectable signal. The responsive transcriptional element may be a Gi responsive transcriptional element, a Gq responsive transcriptional element, or a Gi/Gq responsive transcriptional element. The test compound is contacted to the cell surface under conditions that permit binding of a GPR40 ligand to the GPR40 receptor, and reporter gene expression is detected.

25 A further aspect of the present invention is a method of screening a compound for GPR40 antagonist activity. The method comprises providing a cell expressing on its surface a GPR40 receptor, and containing a reporter construct comprising a responsive transcriptional element and a reporter gene, where expression of the reporter gene produces a detectable signal. The responsive transcriptional element may be a Gi  
30 responsive transcriptional element, a Gq responsive transcriptional element, or a Gi/Gq

responsive transcriptional element. The cell surface is exposed to a test compound and a GPR40 agonist under conditions that permit binding of a GPR40 ligand to the GPR40 receptor, and reporter gene expression is detected. Decreased reporter gene expression in the presence of both test compound and agonist, compared to reporter gene expression in the presence of the GPR40 agonist only, indicates that the test compound is a GPR40 antagonist.

A further aspect of the present invention is a method of screening a compound for GPR40 antagonist activity where the method comprises detecting whether a test compound decreases glucose-stimulated insulin release from, or production by, mammalian pancreatic beta cells in the presence of a GPR40 agonist, compared to glucose-stimulated insulin release that would occur due to the presence of the GPR40 agonist.

A further aspect of the present invention is a method of screening a compound for GPR40 agonist activity, comprising detecting whether the compound binds to GPR40 and increases glucose-stimulated insulin release from, or production by, mammalian pancreatic beta cells.

### Brief Description of the Figures

**Figure 1** shows the calcium response (measured by change in fluorescence) of GPR40 to the fatty acid elaidic acid in HEK-293 cells transiently expressing either GPR40, the 7TM receptor GPRW, or the 7TM receptor HLTEX11.

**Figure 2** shows the results of assays using transiently transfected melanophores to determine the G-protein coupling of GPR40. Increasing amounts of transfected GPR40 DNA produced an increase in  $G\alpha_i/G\alpha_o$  activity, indicating that constitutive activity of GPR40 is through  $G\alpha_i$ .

**Figure 3a** graphs the fold-response of CHO cells expressing CRE-luciferase human GPR40; there was no increase in reporter activity detected in the CRE system at any dose of GPR40 expression vector.

**Figure 3b** graphs the fold-response of CHO cells expressing CRE-luciferase Human Gs-coupled receptor; there was significant increase in luciferase activity detected

with increasing doses of expression vector.

**Figure 3c** graphs the effect of calcitonin and eicosatriynoic acid on luciferase activity from CHO cells expressing CRE-luciferase human GPR40 or CHO cells expressing CRE-luciferase with no receptor. Luciferase production was increased by calcitonin in both the cells containing reporter only and cells containing GPR40, whereas no significant effect of 5,8,11 Eicosatriynoic acid was detected at any concentration tested.

**Figure 4A** shows changes in L-alpha-Lysophosphatidic acid (LPA) stimulated reporter activity, with and without pertussis toxin (PTX) treatment. Host cells consisted of CHO cells containing GAL4/Elk1-luciferase reporter constructs; 3G8 and 5F6 are clonal CHO cell lines transfected with human GPR40 and containing GAL4/Elk1-luciferase reporter construct. Luciferase production in host cells due to LPA-stimulation was abolished by pertussus toxin treatment.

**Figure 4B** shows changes in 5,8,11-Eicosatriynoic acid stimulated reporter activity, with and without pertussis toxin (PTX) treatment. Cells are as described for Figure 4A. Luciferase production due to 5,8,11-Eicosatriynoic acid stimulation was increased in cells transfected with GPR40, and was not inhibited by pertussis toxin treatment.

**Figure 4C** shows changes in thrombin stimulated reporter activity, with and without pertussis toxin (PTX) treatment. Cells are as described for Figure 4A. Luciferase production due to thrombin stimulation was only partially attenuated following pertussis toxin treatment.

**Figure 4D** shows the percent inhibition of luciferase activity following pertussin toxin treatment in CHO cells expressing GAL4/ELK1-luciferase only (host cell), and five different CHO cell clones expressing GAL4/ELK1-luciferase and human GPR40.

**Figure 4E** shows basal luciferase counts from CHO cells expressing GAL4/Elk1-luciferase only (host cell), and different CHO cell clones expressing GAL4/ELK1-luciferase and human GPR40, either with or without pertussis toxin treatment. Clones transfected with GPR40 expression construct and ELK-Gal4 reporter exhibited an increased basal reporter activity compared to that of the host cell line, but this was

decreased by up to 70% following pertussis toxin treatment.

### Detailed Description

GPR40 is a recently described orphan G-protein coupled receptor isolated during a  
5 search for novel receptors (Sawzdargo, M. (1997) Biochem. Biophys. Res. Commun. 239  
(2), 543-547). The polynucleotide sequence of human GPR40 (GenBank Accession no.  
NM\_005303) is given in SEQ ID NO:1 herein, and the amino acid sequence of the  
polypeptide encoded thereby is given in SEQ ID NO:2 herein.

More recently a mouse GPR40 orthologue has been cloned and sequenced (US  
10 patent application no. 60/234,253, SmithKline Beecham). The polynucleotide sequence  
of mouse GPR40 is given in SEQ ID NO:3 herein and the polypeptide encoded thereby  
has the amino acid sequence of SEQ ID NO:4 herein.

The present invention is based on the unusual and unexpected finding that the  
human GPR40 receptor is specifically activated by both saturated and unsaturated fatty  
15 acids. Further, it has been shown that GPR40 is expressed in the human pancreas, more  
specifically the  $\beta$  cells of the pancreatic islets.

The present inventors determined that the GPR40 receptor can constitutively act  
through a pertussis toxin sensitive G protein. Both  $G_i$  and  $G_o$  (and subtypes thereof)  
proteins are known to be pertussis toxin sensitive. However, the present inventors  
20 carried out further research to determine that fatty acids were GPR40 ligands that acted  
via  $G_q$  protein (pertussis toxin insensitive G protein).

Many studies have shown that acute treatment of islets with fatty acids (1-3hrs)  
stimulates glucose-stimulated insulin secretion, whereas longer periods of treatment (6-  
24hrs) are inhibitory to glucose-stimulated insulin secretion (Warnotte C. et al (1994).  
25 Diabetes 43(5):703-711; Zhou YP. et al (1996) Metabolism: Clinical & Experimental.  
45(8):981-986). There is mounting evidence that shows that an increase in plasma fatty  
acid levels may be detrimental to  $\beta$ -cell function (lipotoxicity). Studies in the ZDF rats,  
frequently used as a model of diabetes, have shown that increased levels of circulating  
fatty acids leads to the production of ceramide, activation of nitric oxide synthase, and  
30 apoptosis of  $\beta$ -cells (Shimabukuro M., et al (1998) Proc. Natl. Acad. Sci. U.S.A. 95:

2498-2502). We have found that expression of GPR40 is enriched in human pancreatic islets by 2-100 fold (four individual samples analysed) for an equivalent RNA input compared to human total pancreatic RNA. This tissue localisation suggests that the receptor may have a role in  $\beta$ -cell function. The finding that fatty acids are ligands for GPR40 may aid in the elucidation of GPR40 function in the  $\beta$ -cell. A consequence of acute receptor activation may therefore be enhancement of glucose-stimulated insulin secretion whereas longer periods may be expected to lead to decreases in glucose-stimulated insulin secretion and genes associated with glucose-sensitivity of  $\beta$ -cells e.g. glucokinase. On this basis, it is possible that either agonists or antagonists to GPR40 may be of therapeutic value for diabetes, and particularly type 2 diabetes. Treatment with a GPR40 agonist may be expected therefore to increase glucose-stimulated insulin secretion from  $\beta$ -cells in the short term. GPR40 antagonists may be of therapeutic value with respect to possible reductions in lipotoxicity and thus function to improve  $\beta$ -cell function.

This identification of fatty acids as ligands for GPR40 therefore facilitates the development of screening methods for identifying agonists and antagonists of the receptor.

Accordingly, the present invention provides a method of identifying compounds which bind to and activate (agonist) or inhibit activation (antagonist) of the GPR40 receptor which method comprises using GPR40, in combination with fatty acid GPR40 ligands of C6-C23 saturated or unsaturated fatty acid, optionally containing up to 6 alkene or 3 acetylene bonds, optionally cyclic or branched, and optionally substituted with 1-3 hydroxy groups.

As used herein, "GPR40" refers to a receptor polypeptide having at least 95% identity to the polypeptide sequence given in SEQ ID NO:2, and having GPR40 function; GPR40 receptor polypeptides used in the methods of the present invention are preferably mammalian, and more preferably human. GPR40 also refers to derivatives of the receptor useful in the screening or rational drug design methods disclosed herein. Such derivatives include portions of the polypeptide of SEQ ID NO:2 which retain ligand binding function.

Preferably the fatty acid ligand is selected from the group consisting of:

- 1) trans-retinoic acid (Vitamin A acid; Tretinoin)  $C_{20}H_{28}O_2$  (Unsat)
- 2) cis-4,7,10,13,16,19-Docosahexaenoic acid  $C_{22}H_{32}O_2$  (Unsat)
- 3) Palmitic acid  $C_{16}H_{32}O_2$  (Sat)
- 4) Pentadeconoic acid  $C_{15}H_{30}O_2$  (Sat)
- 5) Elaidic acid  $C_{18}H_{34}O_2$  (Unsat)
- 6) Petroselinic acid  $C_{18}H_{34}O_2$  (Unsat)
- 7) Heptadecanoic acid  $C_{17}H_{34}O_2$  (Sat)
- 8) Tridecanoic acid  $C_{13}H_{26}O_2$  (Sat)
- 9) Lauric acid (Dodecanoic acid)  $C_{12}H_{24}O_2$  (Sat)
- 10) Arachidonic acid  $C_{20}H_{32}O_2$  (Unsat)
- 11) Linolenic acid  $C_{18}H_{30}O_2$  (Unsat)
- 12) Palmitoleic acid  $C_{16}H_{30}O_2$  (Unsat)
- 13) Capric acid  $C_{10}H_{20}O_2$  (Sat)
- 14) Myristic acid  $C_{14}H_{28}O_2$  (Sat)
- 15) Stearic acid  $C_{18}H_{36}O_2$  (Sat)
- 16) Undecanoic acid  $C_{11}H_{22}O_2$  (Sat)
- 16) Eicosatriynoic acid

In this list "sat" refers to saturated and "unsat" refers to unstaured fatty acid.

- 5       As used hereinbelow the term "fatty acid ligand" is defined as a fatty acid ligand of the invention, which ligand is a C6-C23 saturated or unsaturated fatty acid, optionally containing up to 6 alkene or 3 acetylene bonds, optionally cyclic or branched, and optionally substituted with 1-3 hydroxy groups.

- 10       Compounds identified using the methods of the invention may be useful in the treatment of type II diabetes (non-insulin dependent diabetes mellitus, NIDDM) and obesity, glucose intolerance, insulin resistance, neurodegenerative disease (for example Alzheimer's disease) and other indications such as stroke.

- 15       In a first embodiment, an agonist or antagonist of GPR40 may be identified by contacting a cell expressing on the surface thereof the receptor GPR40, said receptor being associated with a second component capable of providing a detectable signal in

response to the binding of a compound to said receptor, with a compound to be screened under conditions to permit binding to the receptor; and determining whether the compound binds to, and activates, or inhibits, the receptor, by detecting the presence or absence of a signal generated from the interaction of the compound with the receptor, optionally in the presence of labeled or unlabeled ligand, for example a fatty acid ligand.

In a further embodiment, an agonist or antagonist of GPR40 may be identified by determining the inhibition of binding of a ligand (e.g., a fatty acid ligand) to cells which have the GPR40 receptor on the surface thereof, or to cell membranes containing the receptor, in the presence of a candidate compound and under conditions permitting binding to the receptor. By determining the amount of ligand bound to the receptor, a compound capable of causing reduction of binding of a ligand is identified as an agonist or antagonist of GPR40.

In general, such screening methods involve providing appropriate cells which express GPR40 on the surface thereof. Such cells include cells from mammals (e.g., Chinese hamster ovary (CHO), HEK (human embryonic kidney), Beta cells of pancreatic islets), melanophore cells, and cells of *Drosophila* or *E. coli*. In particular, a polynucleotide encoding GPR40 is employed to transfect cells to thereby express said receptor. Construction of expression vectors comprising a GPR40-encoding polynucleotide and transfection of cells with said GPR40 expression vectors can be achieved using standard methods, as described in, for example, Sambrook *et al.*, Molecular Cloning: A Laboratory Manual, 2<sup>nd</sup> Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989). Receptor expression may be transient or stable. Preferably, the expression is stable. More preferably a mammalian cell line is transfected with an expression vector comprising a nucleic acid sequence encoding the GPR40 receptor, for example the polynucleotide of SEQ ID NO:1, or the coding region thereof, and the cell line then cultured in a culture medium such that the receptor is stably expressed on the surface of the cell. The expressed receptor is then contacted with a test compound to observe binding, stimulation or inhibition of a functional response, in the presence or absence of a ligand (such as a fatty acid ligand as hereinabove described).

Assays as described herein may utilize intact cells expressing functional GPR40,

or cell membranes containing the receptor, as is known in the art.

Alternatively a soluble portion of the GPR40 receptor (ie. not membrane-bound) comprising the ligand binding domain may be expressed in the soluble fraction, either in the intracellular compartment or secreted out of the cell into the medium. Techniques for  
5 the isolation and purification of expressed soluble receptors are well known in the art.

An example of a screening method includes expressing GPR40 in which the receptor is linked to phospholipase C or D. Representative examples of such cells include, but are not limited to, endothelial cells, smooth muscle cells, and embryonic  
10 kidney cells. The screening may be accomplished as hereinabove described by detecting activation of the receptor or inhibition of activation of the receptor from the phospholipase second signal.

Another method involves screening for compounds which are ligands of GPR40 by determining the inhibition of binding of a labeled ligand (such as a fatty acid ligand) to  
15 cells which have the receptor on the surface thereof, or to cell membranes containing the receptor. Such a method involves transfecting a eukaryotic cell with DNA encoding GPR40 such that the cell expresses the receptor on its surface. The cell is then contacted with a test compound in the presence of a labeled form of a ligand, such as a labelled fatty acid ligand. The ligand can be labeled using any suitable method as is known in the art,  
20 including radioactivity (e.g., with  $^{125}\text{I}$ iodine), fluorescent compounds,  $\beta$ -glucuronidase, luciferase, etc. The amount of labeled ligand bound to the receptors is measured, e.g., by measuring radioactivity associated with transfected cells or membranes from these cells. If the test compound binds to the receptor, the binding of labeled ligand to the receptor is inhibited as determined by a reduction of labeled ligand which binds to the receptors.  
25 This method is generally referred to as a binding assay.

A further screening method involves the use of mammalian cells which are transfected to express functional GPR40. The cells are loaded with an indicator dye that produces a fluorescent signal when bound to calcium, and the cells are contacted with a test substance and/or a receptor agonist, such as a fatty acid ligand. Any change in  
30 fluorescent signal is measured over a defined period of time using, for example, a

fluorescence spectrophotometer or a fluorescence imaging plate reader. A change in the fluorescence signal pattern generated by the ligand indicates that a compound is a potential agonist for the receptor. A decrease in the fluorescence signal pattern generated by the agonist in the presence of the test compound indicates that a compound is potential  
5 antagonist for the receptor.

Another method involves screening for GPR40 ligands by determining effects on receptor-mediated cAMP and/or adenylyl cyclase accumulation. Such a method involves transiently or stably transfecting a eukaryotic cell with GPR40 to express the receptor on the cell surface. The cell is then exposed to test compounds in the presence or absence of  
10 a GPR40 agonist, such as a fatty acid. The change in cAMP level is then measured over a defined period of time, for example, by radio-immuno or protein binding assays (for example using FlashPlates (PerkinElmer Life Sciences) or a scintillation proximity assay). Changes in cAMP levels can also be determined by directly measuring the activity of the enzyme, adenylyl cyclase, in broken cell preparations. If the test  
15 compound binds the receptor and affects GPR40 activation, the levels of GPR40-mediated cAMP, or adenylyl cyclase activity, will be reduced or increased.

Various methods are known in the art to detect GPCR activation, depending in part on the G-protein to which the receptor is coupled. For example, activation of the Gi pathway decreases adenylyl cyclase activity, and Gi signaling may be detected by  
20 measuring the inhibition of forskolin-stimulated cAMP accumulation. (Wong et al., Methods Enzymology, 238:81 (1994)).

The use of reporter-based assays is known in the art to identify GPCR ligands. In general, such assays utilize recombinant cells expressing a functional target receptor protein whose signal transduction activity is modulated by interaction with an  
25 extracellular ligand, with the transduction activity being able to generate a detectable signal. See, e.g., Chen et al., J. Pharmacol. Toxicol. 42:199 (1999); Stables et al., J. Recept. Signal Transduct. Res. 19:395 (1999).

To illustrate, the transduced intracellular signal is initiated by the specific  
30 interaction of an extracellular ligand with the receptor on the cell surface. This interaction

initiates a series of intracellular events that result in a rapid and detectable change in the transcription or translation of a gene. By selecting transcriptional regulatory sequences that are responsive to the transduced intracellular signals and operatively linking the selected promoters to reporter genes whose transcription, translation or ultimate activity is readily detectable and measurable, the transcription based assay indicates whether a specific receptor interacts with a test compound to influence intracellular transduction. See, e.g., Himmler et al., J. Receptor Res. 13:79 (1993); Weyer et al., Receptors and Channels 1:193 (1993); Bevan et al., Neuroreport 9:2703 (1998).

Using such assays, test compounds that induce receptor signaling can be identified (agonists). Alternatively, the assay can test compounds for the ability to potentiate the induction response generated by treatment of the cell with a known agonist. If the test compound does not appear to directly induce the activity of the receptor protein, the assay may be repeated with the recombinant cell first being contacted with a known activator of the target receptor to induce the signal transduction pathway (and generate the detectable signal). Test compounds may then be assayed for the ability to antagonize (inhibit or block) the activation of the receptor by a known agonist activator.

As used herein, an "agonist" refers to agents which either induce activation of receptor signalling pathways, e.g., such as by mimicking a ligand for the receptor, as well as agents which potentiate the sensitivity of the receptor to a ligand, e.g., lower the concentrations of ligand required to induce a particular level of receptor-dependent signalling.

Cells utilized in such assays may express the receptor of interest endogenously, or may be engineered to express a heterologous target receptor protein. Methods for introducing heterologous DNA into eukaryotic cells are known in the art and any suitable method may be used. As is known in the art, it may be desirable to inactivate one or more endogenous genes of the host cells (e.g., the gene for the homologous receptor). Additionally, other proteins involved in signal transduction from the target receptor can be inactivated, or complemented with an ortholog or paralog from another organism, e.g., yeast G protein subunits can be complemented by mammalian G protein subunits in yeast cells also engineered to express a mammalian G protein coupled receptor.

The host cell may contain a reporter construct comprising a reporter gene in operative linkage with one or more transcriptional regulatory elements responsive to the signal transduction activity of the receptor protein. Exemplary reporter genes include enzymes (e.g., luciferase, phosphatase, secreted alkaline phosphatase,  $\beta$ -galactosidase, green fluorescent protein, beta lactamase, chloramphenicol acetyl transferase) which can produce a spectrometrically active label (e.g., changes in color, fluorescence or luminescence), or a gene product which alters a cellular phenotype (e.g., cell growth, drug resistance or auxotrophy).

As used herein, a "reporter gene construct" is a nucleic acid molecule that includes a reporter gene operatively linked to transcriptional regulatory sequence(s) or element(s) which control transcription of the reporter gene. The activity of at least one or more of these control sequences is directly or indirectly regulated by the target receptor protein. The transcriptional regulatory sequences include the promoter and other regulatory regions, such as enhancer sequences, that modulate the activity of the promoter, or regulatory sequences that modulate the activity or efficiency of the RNA polymerase that recognizes the promoter, or regulatory sequences are recognized by effector molecules, including those that are specifically induced by interaction of an extracellular signal with the target receptor. For example, modulation of the activity of the promoter may be effected by altering the RNA polymerase binding to the promoter region, or, alternatively, by interfering with initiation of transcription or elongation of the mRNA. Such sequences are herein collectively referred to as transcriptional regulatory elements or sequences. In addition, the construct may include sequences of nucleotides that alter translation of the resulting mRNA, thereby altering the amount of reporter gene product.

The level of expression of the reporter gene provides the receptor-dependent detection signal. The amount of transcription from the reporter gene may be measured using any suitable method known to those of skill in the art. For example, specific mRNA expression may be detected using Northern blots or real-time quantitative Polymerase Chain Reaction (PCR) such as that available as TAQMAN<sup>®</sup>; or specific protein product may be identified by a characteristic stain or an intrinsic activity. The reporter gene may encode a gene product that, by enzymatic activity, gives rise to a

detection signal based on color, fluorescence, or luminescence. The response of the reporter gene is then compared to the response in either the same cell in the absence of the test compound, or it may be compared with the response in a substantially identical cell that lacks the specific receptors. Any statistically or otherwise significant difference in the response indicates that the test compound has altered the activity of the specific receptor.

Agonists and antagonists are "receptor effector" molecules that modulate signal transduction via the receptor. Receptor effector molecules are capable of binding to the receptor, though not necessarily at the binding site of the natural ligand. Receptor effectors can modulate signal transduction when used alone (surrogate ligands), or can alter signal transduction in the presence of the natural ligand, either to enhance or inhibit signaling by the natural ligand. For example, "antagonists" are molecules that block or decrease the signal transduction activity of receptor, e.g., they can competitively, noncompetitively, and/or allosterically inhibit signal transduction from the receptor, whereas "agonists" potentiate, induce or otherwise enhance the signal transduction activity of a receptor. The terms "receptor activator" and "surrogate ligand" refer to an agonist which induces signal transduction from a receptor.

"Signal transduction" is the processing of chemical signals from the cellular environment through the cell membrane, and may occur through one or more of several mechanisms, such as phosphorylation, activation of ion channels, effector enzyme activation via guanine nucleotide binding protein intermediates, formation of inositol phosphate, activation of adenylyl cyclase, and/or direct activation (or inhibition) of a transcriptional factor. The term "modulation of a signal transduction activity of a receptor protein" in its various grammatical forms, as used herein, designates induction and/or potentiation, as well as inhibition of one or more signal transduction pathways downstream of a receptor.

Methods utilizing pigment cells for identifying GPCR ligands are known in the art. In general, the method utilizes pigment cells capable of dispersing or aggregating pigment in response to a receptor-mediated stimulus, and which express an exogenous G-protein coupled receptor. The condition of the pigment (aggregation or dispersion) in

response to a test compound indicates whether the test compound interacts with the expressed GPCR. See, e.g., US Patent No. 5,462,856; US Patent No. 6,051,386.

*Xenopus laevis* melanophores are able to functionally express recombinant receptors which couple via G-proteins to adenylate cyclase or phospholipase C (PLC).

5 Receptor-mediated stimulation of either of these enzymes causes dispersion of melanosomes, while receptor stimulation that inhibits adenylate cyclase induces melanosome aggregation. Various assays have been devised utilizing *Xenopus* melanophores transfected with a GPCR of interest, to detect the agonist (or antagonist) effects of test compounds on the receptors. See, e.g., Chen et al., *J. Pharmacol. Toxicol.* 42:199 (1999); McClintock et al., *Anal. Biochem.* 209:298 (1993); Potenza et al., *Anal*  
10 *Biochem* 206:315 (1992); Potenza et al., *Pigment Cell Res* 5:372 (1992)).

The Fluorometric Imaging Plate Reader (FLIPR™, Molecular Devices, UK) measures intracellular fluorescence in multi-well microplates. Where a GPCR couples to  
15 a calcium signalling pathway, the FLIPR system can be used with cells expressing the G-protein coupled receptor and containing calcium-sensitive fluorescent dyes, to monitor the intracellular calcium flux that occurs after activation of the G-coupled receptor. See, e.g., Kirk et al., *J. Biomolecular Screening*, 1:75 (1996).

Certain GPCRs do not couple with calcium signalling pathways, while others  
20 may not signal through calcium pathways depending on the type of cell in which the receptor is expressed. In such situations, chimeric G proteins have been utilized to link the receptor to calcium signalling in transfected cells, thereby allowing the use of high throughput FLIPR assays. (Wada et al., *Developing a High Throughput Functional Screening Assay for Gi/o-coupled Cloned Receptors using the FLIPR™ System*,  
25 published by Molecular Devices, 2001; Coward et al., *Chimeric G-proteins allow a high-throughput signaling assay of Gi-coupled receptors*, *Analytical Biochemistry* 270:242 (1999)). Coward et al report the use of chimeric G proteins that allow Gi-coupled receptors to signal through Gq and generate a detectable calcium response in CHO cells, measured using FLIPR™ technology.

Accordingly the present invention provides methods of screening test compounds to determine whether the compound is a GPR40 receptor ligand, by detecting whether the test compound competitively inhibits the binding of a fatty acid GPR40 ligand to a GPR40 receptor. The fatty acid ligand may be labeled with a detectable label as is well known in the art. As used herein, detectable labels include moieties that can be detected by visual inspection (e.g., that include or produce colored elements), or detected with the aid of artificial detection systems, including, e.g., optical systems, spectroscopic systems, radiographic systems, and the like. The detectable label may be selected from among radioisotopes, fluorescent dyes, and enzymes.

The present invention also provides methods of screening test compounds for GPR40 antagonist activity. One such method utilizes a test cell expressing on its surface a GPR40 receptor, and containing a reporter construct comprising a Gq responsive transcriptional element and a reporter gene, where expression of the reporter gene produces a detectable signal. The test cell surface is exposed to a GPR40 fatty acid agonist under conditions that permit binding of the GPR40 fatty acid agonist to the GPR40 receptor, and then the cell surface is exposed to the test compound under similar conditions. The detectable signal produced by the reporter gene is then measured, and compared to the detectable signal that would be expected to occur in the presence of the GPR40 fatty acid agonist only (i.e., in the absence of the test compound). This comparison may be a side-by-side comparison of the detectable signal in a control cell (i.e., a cell similar in all ways to the test cell, but exposed to the GPR40 fatty acid agonist in the absence of test compound). Alternatively the comparison may be to a standard that is determined prior to or after the test cell is processed, i.e., a control experiment that was conducted prior to or after the detectable signal to establish a quantitative or qualitative standard that is used in the comparison.

The present invention further provides methods of screening a compound for GPR40 agonist activity. One such method utilizes a cell expressing GPR40 receptors on its surface, and containing a reporter construct comprising a responsive transcriptional element and a reporter gene, where expression of the reporter gene produces a detectable signal. The transcriptional element may be a Gi responsive transcriptional element, a Gq

responsive transcriptional element, or a Gi/Gq responsive transcriptional element. The cell surface is contacted with (i.e., exposed to) a test compound, under conditions that permit binding of a GPR40 ligand to the GPR40 receptor, and any reporter gene expression is detected (indicating that the test compound has GPR40 agonist activity).

5       The present invention further provides methods of screening a compound for GPR40 antagonist activity. One such method utilizes a cell expressing on its surface a GPR40 receptor, and containing a reporter construct comprising a responsive transcriptional element and a reporter gene, where expression of the reporter gene produces a detectable signal. The responsive transcriptional element may be a Gq  
10 responsive transcriptional element or a Gi/Gq responsive transcriptional element. The cell surface is exposed to (i.e., contacted with) a test compound and a GPR40 agonist, under conditions that permit binding of the GPR40 agonist to the GPR40 receptor. Subsequent reporter gene expression is detected, and decreased reporter gene expression in the presence of both test compound and agonist (compared to reporter gene expression  
15 expected in the presence of said GPR40 agonist only), indicates that the test compound is a GPR40 antagonist. As described previously herein, comparison of the reporter gene expression in the test cell may be made with a control cell, or to a known standard quantitative or qualitative response.

As used herein, a Gq responsive transcriptional element is one that responds (is  
20 activated) upon activation of a  $G\alpha_q$  (or  $G\alpha_q$  subtype) linked receptor. As is known in the art, Gq proteins are pertussis toxin insensitive G-proteins. As used herein, a Gi responsive transcriptional element is one that responds (is activated) upon activation of a  $G\alpha_i$  (or  $G\alpha_i$  subtype) linked receptor. As is known in the art, Gi proteins are pertussis toxin sensitive G-proteins (as are  $G_o$ ). As used herein, a Gq/Gi responsive transcriptional  
25 element is one that responds (is activated) upon activation of a  $G\alpha_q$ -linked receptor (or  $G\alpha_q$  subtypes), or a  $G\alpha_i$ -linked receptor (or  $G\alpha_i$  subtypes). One such Gq/Gi responsive transcriptional element is the GAL4/Elk-1 (see, e.g., Bevan et al., Neuroreport 9:2703 (1998)).

The present invention further provides methods of screening a compound for  
30 GPR40 antagonist activity, by detecting whether the compound decreases glucose-

stimulated insulin release from mammalian pancreatic beta cells (or insulin production within mammalian pancreatic beta cells) in the presence of a GPR40 agonist, compared to glucose-stimulated insulin release that would occur due to the presence of the GPR40 agonist. the "expected" insulin release due to the presence of the GPR40 agonist may be established by carrying out a simultaneous or subsequent control experiment, or may have be a pre-determined standard. Various methods are known in the art for measuring insulin release from, or production within, cells. Any suitable method may be utilized in the present methods.

The present invention further provides a method of screening a compound for GPR40 agonist activity, by detecting whether the compound binds to GPR40 and increases glucose-stimulated insulin release from (or insulin production within) mammalian pancreatic beta cells.

Conversely, further receptors for which the fatty acid ligands of the invention act as ligands may be identified by screening potential candidates against a fatty acid ligand in a suitable assay, for instance determining potential inhibition of forskolin-elevated cAMP levels.

Kits suitable for carrying out the present methods may contain:

(a) GPR40 and one or more labeled or unlabeled fatty acid ligands;  
(b) a recombinant cell expressing GPR40 and one or more labeled or unlabeled fatty acid ligands; or

(c) a cell membrane expressing GPR40 and one or more labeled or unlabeled fatty acid ligands .

It will be appreciated that in any such kit, (a), (b), or (c) may comprise a substantial component.

Agonists and/or antagonists may be identified from a variety of sources, for instance, from cells, cell-free preparations, chemical libraries and natural product mixtures. Such agonists and/or antagonists may be natural or modified substrates, ligands, receptors, enzymes, etc., as the case may be, of any of the fatty acid ligands of the invention; or may be structural or functional mimetics of the polypeptide of the present invention (see Coligan

*et al.*, *Current Protocols in Immunology* 1(2):Chapter 5 (1991)).

Examples of potential antagonists include:

- (a) antibodies or, in some cases, oligonucleotides, which bind to the receptor but do not elicit a second messenger response such that the activity of the receptor is prevented;
- (b) fatty acids which are closely related to a ligand of GPR40 but which elicit no response from the GPR40 receptor;
- © small molecules which bind to GPR40, making it inaccessible to ligands such that normal biological activity is prevented, for example, small peptides or peptide-like molecules;
- (d) soluble forms of GPR40, *e.g.*, fragments of the receptor, which bind to the ligand and prevent the ligand from interacting with membrane bound GPR40.

The invention further provides a method of rational drug design comprising the steps:

- a) probing the structure of a fatty acid ligand-binding site on the GPR40 receptor with said fatty acid ligand or a derivative thereof;
- b) identifying contacting atoms in the binding site of the GPR40 receptor that interact with a fatty acid ligand during binding; and
- c) designing agonist or antagonist compounds that interact with the atoms in the binding site identified in (b) to activate (agonist) or inhibit activation (antagonist) of the receptor.

Conversely, the structure of any of the fatty acids ligands when bound to the ligand binding site on the GPR40 receptor can also be determined, enabling the design of further antagonist compounds. Such antagonists bind to the fatty acid ligand, thereby preventing the binding of the fatty acid ligand to the receptor. Methods of using ligands or their derivatives to probe the structure of the ligand binding sites in receptors, and rational drug design based on this structural information are well known in the art (see for example Boyle, S et al *Bioorganic & Medicinal Chemistry* 1994, 2, 101-113; Beck-Sickinger, A.G et al, *European Journal Of Biochemistry* 1994, 225, 947-958; McWherter, C.A et al. *J.Biol.Chem.* 1997, 272, 11874-11880; Horwell, D.C et al. *International*

*Journal of Peptide & Protein Research* 1996, 48, 522-531; Bednarek, M.A. et al. *Peptides* 1999, 20, 401-409).

Compounds identified using the above screening or rational design methods will be of use in therapy. Accordingly, in a further aspect, the present invention provides a compound identified as an agonist or an antagonist of GPR40 for use in therapy, in particular for treating type II diabetes (non-insulin dependent diabetes mellitus, NIDDM) and obesity, glucose intolerance, insulin resistance, neurodegenerative disease (for example Alzheimer's disease) and other indications such as stroke, among others.

Accordingly, in a further aspect, this invention provides a method of treating an abnormal condition related to an excess of GPR40 activity and/or an excess of a GPR40 ligand (for example a fatty acid ligand). The method comprises administering to a patient in need thereof a GPR40 antagonist as hereinbefore described in an amount effective to block or decrease binding of ligands to the receptor, or to inhibit or decrease a second signal, thereby alleviating the abnormal condition.

In still another approach, expression of the gene encoding endogenous GPR40 polypeptide can be inhibited using expression blocking techniques. Known such techniques involve the use of antisense sequences, either internally generated or externally administered (see, for example, O'Connor, *J Neurochem* (1991) 56:560 in *Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression*, CRC Press, Boca Raton, FL (1988)). Alternatively, oligonucleotides which form triple helices ("triplexes") with the gene can be supplied (see, for example, Lee *et al.*, *Nucleic Acids Res* (1979) 6:3073; Cooney *et al.*, *Science* (1988) 241:456; Dervan *et al.*, *Science* (1991) 251:1360). These oligomers can be administered *per se* or the relevant oligomers can be expressed *in vivo*. Synthetic antisense or triplex oligonucleotides may comprise modified bases or modified backbones. Examples of the latter include methylphosphonate, phosphorothioate or peptide nucleic acid backbones. Such backbones are incorporated in the antisense or triplex oligonucleotide in order to provide protection from degradation by nucleases and are well known in the art. Antisense and triplex molecules synthesised with these or other modified backbones also form part of the present invention.

In addition, expression of the human GPR40 polypeptide may be prevented by

using ribozymes specific to the human GPR40 mRNA sequence. Ribozymes are catalytically active RNAs that can be natural or synthetic (see for example Usman, N, et al., Curr. Opin. Struct. Biol (1996) 6(4), 527-33.) Synthetic ribozymes can be designed to specifically cleave GPR40 mRNAs at selected positions thereby preventing translation of the human GPR40 mRNAs into functional polypeptide. Ribozymes may be synthesised with a natural ribose phosphate backbone and natural bases, as normally found in RNA molecules. Alternatively the ribozymes may be synthesized with non-natural backbones to provide protection from ribonuclease degradation, for example, 2'-O-methyl RNA, and may contain modified bases.

Identification of a ligand for GPR40, such as a fatty acid ligand of the invention, allows for the effective identification of polyclonal or monoclonal antibodies raised against GPR40 which are neutralizing antibodies. Such neutralizing antibodies are of use in therapy, in comparison to non-neutralizing antibodies which are ineffective. Accordingly, in a further aspect, the present invention provides for the use of neutralizing antibodies raised against GPR40 in therapy.

Compounds which are active when given orally can be formulated as liquids, for example syrups, suspensions or emulsions, tablets, capsules and, lozenges. A liquid formulation will generally consist of a suspension or solution of the compound or pharmaceutically acceptable salt in a suitable liquid carrier(s) for example, ethanol, glycerine, non-aqueous solvent (for example polyethylene glycol), oils, or water with a suspending agent, preservative, flavouring or colouring agent. A composition in the form of a tablet can be prepared using any suitable pharmaceutical carrier(s) routinely used for preparing solid formulations. Examples of such carriers include magnesium stearate, starch, lactose, sucrose and cellulose. A composition in the form of a capsule can be prepared using routine encapsulation procedures. For example, pellets containing the active ingredient can be prepared using standard carriers and then filled into a hard gelatin capsule; alternatively, a dispersion or suspension can be prepared using any suitable pharmaceutical carrier(s) (for example aqueous gums, celluloses, silicates or oils) and the dispersion or suspension then filled into a soft gelatin capsule. Typical parenteral compositions consist of a solution or suspension of the compound or pharmaceutically

acceptable salt in a sterile aqueous carrier or parenterally acceptable oil, for example polyethylene glycol, polyvinyl pyrrolidone, lecithin, arachis oil or sesame oil. Alternatively, the solution can be lyophilised and then reconstituted with a suitable solvent just prior to administration. A typical suppository formulation comprises an active compound or a pharmaceutically acceptable salt thereof which is active when administered in this way, with a binding and/or lubricating agent such as polymeric glycols, gelatins or cocoa butter or other low melting vegetable or synthetic waxes or fats. The invention is further described in the following examples which are intended to illustrate the invention without limiting its scope. In order to facilitate understanding of the following examples certain methods and/or terms will be described.

All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

## Examples

### Example 1

#### TAQMAN® analysis of GPR40 expression in Human Tissues

Generation of samples for TAQMAN® mRNA analysis:

TAQMAN® PCR was performed following the procedure published by Sarau H.M. et al, ("Identification, Molecular Cloning, Expression and Characterisation of a Cysteinyl Leukotriene Receptor", Molecular Pharmacology, 1999, 56, 657-663.) TAQMAN® quantitative PCR was conducted to measure GPR40 mRNA using replicate 96-well plates. A 20µl volume of a PCR master mix (containing 2.5µl TAQMAN® buffer, 6µl 25mM MgCl<sub>2</sub>, 0.5µl of 10mM dATP, 0.5µl of 20mM dUTP, 0.5µl of 10mM dCTP, 0.5µl of 10mM dGTP, 0.25µl Uracil-N-glycosylase, 1µl of 10µM forward primer, 1µl of 10µM reverse primer, 0.5µl 5µM TAQMAN® probe, 0.125µl TaqGold [PE Biosystems], 6.625µl water) was added to each well using Biomek robotics (Beckman Coulter, High

Wycombe, UK), and the plate capped using optical caps (PE Biosystems). The PCR reaction was carried out on an ABI7700 Sequence Detector (PE Biosystems) using the PCR parameters: 50°C for 2 minutes, 95°C for 10 minutes and 45 cycles of 94°C for 15 seconds, 60°C for 1 minute, and the level of mRNA-derived cDNA in each sample was calculated from the TAQMAN® signal using plasmid/genomic DNA calibration standards included in each run. The level of genomic DNA contaminating the original RNA samples was shown to be negligible (<10 copies genomic DNA/ng RNA) by TAQMAN® measurement of genomic sequence for ten genes in replicate samples taken through the reverse transcription procedure described with the omission of reverse transcriptase. Gene-specific reagents for GPR40:

forward primer: 5' GTGGTGCTTAATCCGCTGGT (SEQ ID NO:5)  
reverse primer: 5' TGGCGTTACTTCTGGGACTTG (SEQ ID NO:6)  
probe: 5' CTTGCGTTCTTGCCGCACACACTGT (SEQ ID NO:7)

The results show that the highest levels of GPR40 are expressed in the pancreas; results are shown in Table 1

**Table 1**

GPR40 Body tissue mRNA levels

units: copies of GPR40 mRNA detected/ng mRNA pool

	mean mRNA copies	+/-SEM
Brain	111	12
Pituitary	46	18
Heart	91	47
Lung	30	15

Liver	28	11
Foetal Liver	59	28
Kidney	24	13
Muscle	44	10
Stomach	22	7
intestine	34	21
Spleen	61	4
Lymphocytes	62	24
Macrophage	7	7
adipose	30	6
Pancreas	225	44
Prostate	78	32
Placenta	31	11
Cartilage	5	6
Bone	30	11
Bone marrow	20	12

### Example 2

#### Expression of GPR40 in human pancreatic islet cells

- 5 TAQMAN® analysis of human pancreatic islet RNA vs total pancreatic RNA was carried out as described in Example 1. For an equivalent RNA sample from islet cells and total pancreas, in 3 independent islet samples there was a 2 to 100 fold higher expression of GPR40 than in total pancreas. (Results not shown).

10

### Example 3

#### Expression of GPR40 in MIN6 and BRIN insulinoma cells

#### TAQMAN® Conditions for mouse and rat GPR40 Expression:

- 15 TAQMAN® was performed using 50ng cDNA per reaction with 0.3μM forward primer, 0.3μM reverse Primer, 0.1μM probe and 1x TAQMAN® Master Mix (PE Applied Biosystems, UK). The cycle conditions were as follows 50°C for 2 min, 95°C for 10 min followed by 40 cycles of 95°C for 15 secs, 60°C for 1 min.

**Primer/Probe sets for mouse/rat GPR40**

Forward Primer: 5' AGTTCCTGGGCATCAACATA3' (SEQ ID NO:8)

Reverse Primer: 5' CAAGGGCAGAAAGAAGAGCAGA3' (SEQ ID NO:9)

5 Probe: 5' AATGGCTCCCCGGTCTGCCTGG3' (SEQ ID NO:10)

Results are shown in **Table 2**. Results show that GPR40 was expressed in both insulinoma cell-lines, confirming  $\beta$ -cell localisation. The results further showed that expression of GPR40 in MIN6 cells is much greater than in BRIN cells.

**Table 2**

Insulinoma cell line	Av Ct value <sup>1</sup>	Relative value (mouse std curve) normalised (HPRT <sup>2</sup> )	Relative value (rat std curve) normalised (HPRT <sup>2</sup> )
MIN6 (Mouse Insulinoma )	22.7	0.95	119946.9
BRIN (rat insulinoma)	37.1	$3.23 \times 10^{-5}$	0.09

10 1) Ct value refers to the cycle number of TAQMAN® necessary to detect gene expression for that transcript.

2) HPRT = Hypoxanthine-Guanine Phosphoribosyltransferase

**Example 4**

15 **GPR40 expression is increased in *ob/ob* mouse pancreas**

TAQMAN® analysis was carried out to determine GPR40 levels in C57B1/6 lean mice compared with C57B1/6 *ob/ob* homozygous knock-out mice. These levels were further compared with levels of insulin mRNA. The results are shown in **Table 3**.

**Table 3**

	Relative Quantity GPR40 <sup>1</sup> (mRNA $\pm$ s.e.m.)	Relative Quantity Insulin <sup>1</sup> (mRNA $\pm$ s.e.m.)
C57/B1/6 lean mice (n=10)	12 $\pm$ 28	78 $\pm$ 7

C57B1/6 <i>ob/ob</i> mice (n=10)	113 $\pm$ 23	960 $\pm$ 145
-------------------------------------	--------------	---------------

1) relative measure from a standard curve created using known amounts of cDNA made from total RNA with random primers.

### Example 5

#### GPR40 and fatty acids – receptor ligand pairing

Cells expressing the GPR40 receptor exhibit a dose-related increase in intracellular calcium levels in response to fatty acid ligands, as measured by FLIPR technology. **Figure 1** shows the FLIPR response detected following addition of elaidic acid (trans oleic acid, 9-octadecanoic acid) to HEK-293 cells transiently expressing the GPR40 receptor. No response was detected in cells transiently expressing two other 7TM receptors (PGRW and HLTEX11). The assays utilized a G-protein cocktail (+GPC) that contained chimeric and promiscuous G-proteins, as is known in the art of screening orphan G-protein coupled receptors.

GPR40 stimulation by other fatty acid ligands, using FLIPR technology as described herein, is shown in **Table 4** below (pEC50 =  $-\log$  EC50 where EC50 is 'effective concentration'; EC50 is half of the concentration of compound necessary for a maximal response)

**Table 4**

	Mean pEC50	Mean pEC50	Mean pEC50		Mean pEC50 of 3 expts	(+/-) sem
<b>Unsaturated Fatty Acids</b>						
Palmitoleic acid C <sub>16</sub> H <sub>30</sub> O <sub>2</sub> (9Z)	4.82	4.81	4.96		4.86	0.05
Linolenic acid C <sub>18</sub> H <sub>30</sub> O <sub>2</sub> (9Z, 12Z, 15Z)	4.93	4.66	5.12		4.90	0.13
Linoleic acid C <sub>18</sub> H <sub>32</sub> O <sub>2</sub> (9Z, 12Z)	4.78	4.68			4.73	n=2
12. Elaidic acid C <sub>18</sub> H <sub>34</sub> O <sub>2</sub> (9E)	5.03	5.27	5.18		5.16	0.07
Oleic acid C <sub>18</sub> H <sub>34</sub> O <sub>2</sub> (9Z)	4.55	4.23			4.39	n=2
Petroselinic acid C <sub>18</sub> H <sub>34</sub> O <sub>2</sub> (6Z)	4.88*	4.97	5.03		5.00	0.02

All-trans-retinal (Vitamin A aldehyde) C <sub>20</sub> H <sub>28</sub> O	4.08	4.23			4.16	n=2
All-trans-retinoic acid (Vitamin A acid; Tretinoin) C <sub>20</sub> H <sub>28</sub> O <sub>2</sub>	5.56	5.52	5.65		5.58	0.04
9-cis-retinoic acid (9-cis-tretinoin) C <sub>20</sub> H <sub>28</sub> O <sub>2</sub>	4.27	4.52			4.40	n=2
All-trans-retinol (Vitamin A) C <sub>20</sub> H <sub>30</sub> O	No Fit	No Fit				
Arachidonic acid C <sub>20</sub> H <sub>32</sub> O <sub>2</sub> (5Z, 8Z, 11Z, 14Z)	4.74	4.89	5.14		4.92	0.12
cis-4,7,10,13,16,19-Docosahexaenoic acid C <sub>22</sub> H <sub>32</sub> O <sub>2</sub>	5.18	5.53	5.4		5.37	0.10
Erucic acid C <sub>22</sub> H <sub>42</sub> O <sub>2</sub> (13Z)	4.33*	No Fit			4.33	n=1
Nervonic acid C <sub>24</sub> H <sub>46</sub> O <sub>2</sub> (15Z)	No Fit	No Fit				
<b>Saturated Fatty Acids</b>						
n-caproic acid C <sub>6</sub> H <sub>12</sub> O <sub>2</sub>	4.33*	No Fit		6	4.33	n=1
Heptanoic acid C <sub>7</sub> H <sub>14</sub> O <sub>2</sub>	4.33	4.22*		7	4.28	n=2
Caprylic acid C <sub>8</sub> H <sub>16</sub> O <sub>2</sub>	4.25	4.58		8	4.42	n=2
Nonanoic acid C <sub>9</sub> H <sub>18</sub> O <sub>2</sub>	4.16	4.63		9	4.40	n=2
Capric acid C <sub>10</sub> H <sub>20</sub> O <sub>2</sub>	4.74	4.96	4.84	10	4.85	0.06
Undecanoic acid C <sub>11</sub> H <sub>22</sub> O <sub>2</sub>	4.6	4.66	4.83	11	4.70	0.07
Lauric acid (Dodecanoic acid) C <sub>12</sub> H <sub>24</sub> O <sub>2</sub>	4.75	4.93	5.07	12	4.92	0.09
Tridecanoic acid C <sub>13</sub> H <sub>26</sub> O <sub>2</sub>	4.8	4.95	5.05	13	4.93	0.07
Myristic acid C <sub>14</sub> H <sub>28</sub> O <sub>2</sub>	4.95	4.52	5.05	14	4.84	0.16
Pentadecanoic acid C <sub>15</sub> H <sub>30</sub> O <sub>2</sub>	4.97	5.36	5.21	15	5.18	0.11
Palmitic acid C <sub>16</sub> H <sub>32</sub> O <sub>2</sub>	5.08	5.51	5.3	16	5.30	0.12
Heptadecanoic acid C <sub>17</sub> H <sub>34</sub> O <sub>2</sub>	4.86	4.99	5.11	17	4.99	0.07
Stearic acid C <sub>18</sub> H <sub>36</sub> O <sub>2</sub>	4.7	4.78	4.85	18	4.78	0.04
Nonadecanoic acid C <sub>19</sub> H <sub>38</sub> O <sub>2</sub>	4.58	4.46*		19	4.52	n=2
Arachidic acid C <sub>20</sub> H <sub>40</sub> O <sub>2</sub>	No Fit	4.21*		20	4.21	n=1
Heneicosanoic acid C <sub>21</sub> H <sub>42</sub> O <sub>2</sub>	4.45	4.53*		21	4.49	n=2
Behenic acid (Docosanoic acid) C <sub>22</sub> H <sub>44</sub> O <sub>2</sub>	4.29	4.31		22	4.30	n=2
Tricosanoic acid C <sub>23</sub> H <sub>46</sub> O <sub>2</sub>	4.31*	No Fit		23	4.31	n=1
Lignoceric acid C <sub>24</sub> H <sub>48</sub> O <sub>2</sub>	No Fit	No Fit		24	0	

Additional Fatty Acid Compounds	EC50 (M)	EC50 uM	pEC50		
Mean acid (10:3,n-9)	2.51E-06	2.5	5.60		
Linoleic acid (18:2,n-6)	9.54E-06	9.5	5.02		
Linolenic acid (18:3,n-3)	5.50E-06	5.5	5.26		
Gamma-Linolenic acid (18:3,n-6)	8.91E-06	8.9	5.05		
Elaidic acid (18:1, n-9, E)	1.35E-05	13.5	4.87		
17-Octadecynoic acid (17-ODYA) (18:1 x terminal acetylene	7.58E-06	7.6	5.12		
Eicosa-11Z,14Z-dienoic acid (20:2,n-6)	1.07E-05	10.7	4.97		
Eicosa-5Z,8Z-dienoic acid (20:2,n-12)	7.76E-06	7.8	5.11		
Eicosa-11Z,14Z,17Z-trienoic acid (20:3,n-3)	1.12E-05	11.2	4.95		
Dihomo-gamma-linolenic acid (20:3,n-6)	7.24E-06	7.2	5.14		
5,8,11-Eicosatriynoic acid (20:3 x acetylenes)	1.70E-06	1.7	5.77		
Eicosa-5Z,8Z,11Z,14Z,17Z-pentaenoic acid (20:5, n-3)	6.76E-06	6.8	5.17		
Arachidonic acid (20:4,n-6)	9.12E-06	9.1	5.04		
14®, 15(S)-DiHETE (20:4 + 2 hydroxyls)	2.34E-05	23.4	4.63		
Docosa-13Z,16Z,19Z-trienoic acid (22:3,n-3)	6.76E-06	6.8	5.17		
Adrenic acid (22:4,n-6)	1.35E-05	13.5	4.87		
Docosa-7Z,10Z,13Z,16Z,19Z-pentaenoic acid (22.5, n-3)	4.68E-06	4.7	5.33		
Docosa-4Z,7Z,10Z,13Z,16Z,19Z-hexaenoic acid (22:6, n-3)	1.62E-06	1.6	5.79		

**Example 5A**  
**Functional Assays**  
**G-protein Coupling**

Most GPCRs show constitutive activity as expression of the GPCR increases. Experiments were performed to determine whether GPR40 couples through  $G\alpha_s$ ,  $G\alpha_q$  or  $G\alpha_i$ , using constitutively active reporters and following agonist activation.

Constitutive GPR40 activity was examined in melanophores. The melanophores were transiently transfected with 10, 20, 40 and 80ug of GPR40/pJG3.6 plasmid DNA by electroporation and plated into 96-well plates. Forty-eight hours post-electroporation, media was aspirated and replaced with L-15 assay buffer, assay buffer + 25nM Melatonin or assay buffer + 100nM alpha-MSH. These reagents were allowed to incubate for 2 hours and the transmittance (T) was read. Percent constitutive activity was calculated using the following calculation:

$$\frac{(T(L15) - T(MSH))}{(T(Melatonin) - T(MSH))} * 100$$

Results show that increasing amounts of transfected GPR40 DNA produced an increase in the  $G\alpha_i/G\alpha_o$  activity, indicating that constitutive activity of GPR40 is through either the  $G\alpha_i/G\alpha_o$  family of G-proteins. (**Figure 2**)

Specific coupling was determined in mammalian cells using transient expression of a CRE-luciferase reporter for  $G\alpha_s$  and stable cell-lines containing Gal4/ELK-1-luciferase reporter for  $G\alpha_q/G\alpha_i$ . Wild-type CHO cells were co-transfected with 3X-CRE-luciferase reporter vector and various amounts of an expression vector for GPR40. The control cells were transfected with reporter vector only. The reporter gene plasmids were produced in house but similar vectors can be purchased from Stratagene (PATHDETECT®). Cells were plated in a 24 well tissue culture plate at  $1 \times 10^4$  cells/well in DMEM/F12 supplemented with 5% FBS. After 72 hours at 37°C, 5% CO<sub>2</sub>, the cells were transfected

using the liposomal transfection reagent sold under the trademark LIPOFECTAMINE® and with the following combinations of plasmids:

A- 3x-CRE-Luc

B- 3x-CRE-Luc + CMV-GPR40

C - 3x-CRE-Luc + CMV-Gs coupled receptor

The concentration of reporter plasmid for all wells was 0.15µg/well. Concentrations of receptor plasmids were 0.5, 1.0, 5.0, 10.0 ng/well. After 24hrs Cells containing 1ng/well of receptor plasmid or no DNA control and CRE-Luc reporter were treated with fatty acid or the calcitonin which is known to act through a Gs-coupled receptor (Chakraborty, Science 251: 1078 (1991)) at concentrations indicated. Luciferase production was assayed after 4 hours of treatment using a luciferase reporter gene assay kit available under the trademark LUCLITE®, used per manufacturers instructions (Packard Bioscience).

As shown in **Figure 3a**, there was no increase in reporter activity in the CRE system at any dose of GPR40 expression vector. In contrast, as shown in **Figure 3b**, expression of a Gs-coupled receptor elicited a significant increase in luciferase activity.

As shown in **Figure 3c**, although calcitonin elicited an increase in luciferase production in both the cells containing reporter only and those containing GPR40, there was no effect of 5,8,11 Eicosatriynoic acid at any concentration tested, indicating that neither GPR40 constitutive nor agonist-activated activity is coupled via Gαs.

To further test the coupling of GPR40 to Gα<sub>i</sub>/Gα<sub>q</sub> a Gal4/Elk-1 reporter was used. A stable, clonal reporter line was created by transfecting CHO cells with a plasmid that contains an expression construct for GAL4/Elk-1 and a GAL4-luciferase reporter construct. A similar reporter system can now be purchased from Stratagene (under the trademark PATHDETECT®). An expression construct for human GPR40 which contains a G418 resistance construct was transfected into the host CHO GAL4/Elk-1 reporter cell line (10ug plasmid DNA pCR3-GPR40) via electroporation. The cell lines were selected for with 500ug/mL G418 until all cells in the control were dead and clones were then picked. The Gal4/Elk-1 reporter can be activated through Gq or Gi. To confirm coupling via Gi, cells were treated with pertussis toxin treatment which ADP-ribosylates Gα<sub>i</sub>

preventing its interaction with receptors ( Bokoch et al., J. Biol. Chem 258:2072 (1983)). Pertussis toxin has no known effect on  $G\alpha_q$ .

Host cells and those transfected with the GPR40 construct were plated in a 96-well tissue culture plate at 2000 cells/well in DMEM/F12 supplemented with 5% FBS. After 48 hours at 37°C at 5% CO<sub>2</sub>, the cells were placed in serum-free DMEM/F12 in the presence or absence of 1ug/ml pertussis toxin. After 16 hours at 37°C, 5% CO<sub>2</sub> the host cells were treated with: lysophosphatidic acid (oleoyl) (LPA) which couples through  $G\alpha_i$  to activate MAP kinase (Hordijk et al. , J. Biol. Chem. 269: 645 (1994)); thrombin which couples through  $G\alpha_i$  and  $G\alpha_q$  (Hung et al. , J. Biol. Chem. 267:20831 (1992)); or 5,8,11-Eicosatriynoic acid (a GPR40 ligand), at the concentrations indicated. The host cells containing the GPR40 construct were stimulated with or 5,8,11-Eicosatriynoic acid at the concentrations indicated. Controls for lysophosphatidic acid contained 0.5% DMSO; controls for 5,8,11-eicosatrienoic acid contained 0.05% ethanol. Luciferase activity was assayed 4 hours after treatment using a luciferase reporter gene assay kit available under the trademark LUCCLITE®, as per manufacturers instructions (Packard Bioscience).

As shown in **Figure 4A**, in the host cells LPA-stimulated reporter activity was completely abolished by the pertussis toxin treatment confirming coupling through  $G\alpha_i$ .

As shown in **Figure 4C**, luciferase production due to thrombin stimulation was only partially attenuated following toxin treatment confirming  $G\alpha_i/G\alpha_q$  coupling.

As shown in **Figure 4D** and **4E**, clones transfected with GPR40 expression construct and ELK-Gal4 reporter exhibited an increase in basal reporter activity versus the host cell line containing reporter alone, which was decreased by up to 70% following pertussis toxin treatment suggesting constitutive  $G\alpha_i$ -coupled receptor activity.

As shown in **Figure 4B**, 5,8,11-Eicosatriynoic acid had no effect on luciferase production in the host cells, but increased luciferase activity compared to basal in cells containing GPR40 in a manner which was not inhibited by pertussis toxin treatment.

These assays demonstrate that constitutive GPR40 receptor activity is coupled to the  $G\alpha_i$  or  $G\alpha_o$  (and related family members) pertussis toxin sensitive G-proteins. Furthermore, fatty acid agonist stimulated GPR40 receptor activation is coupled through a

pertussis-toxin insensitive G-protein, suggesting that agonist-stimulated GPR40 receptor activation is coupled through  $G\alpha_q$ .

### **Example 6**

#### **Functional Assays**

##### **Calcium Assay**

Receptors stably expressed in HEK 293 cells can demonstrate a robust calcium response to agonists with the appropriate rank order and potency. Basal calcium levels in the HEK 293 cells in GPR40-transfected or vector control cells are in the normal 100 nM to 200 nM range. HEK 293 cells expressing recombinant GPR40 receptors are loaded with the fluorescent indicator FLUO-4 and in a single day > 150 selected ligands are evaluated for agonist-induced calcium mobilization. Agonists presenting a transient calcium mobilization are tested in vector control cells to determine if the calcium response is unique to the transfected receptor cells. When a unique agonist-induced response is identified, the response is reproduced in a separate group of cells and then pharmacologically characterized with concentration response curves for the effective and related ligands.

### **Example 7**

#### **Membrane Preparation and Screen for GPR40**

Fatty acid ligand binding competition assays are useful for the discovery of antagonists and agonists of the GPR40 receptor. As a source of the GPR40 receptor, CHO or HEK 293 cells stably transfected with the GPR40 receptor are useful; other cells transfected with the GPR40 receptor or cells that naturally show a high level expression of the GPR40 receptor may also be employed. Typically the culture of cells expressing the GPR40 receptor is scaled up to 30L and cells are recovered by centrifugation at 600 x g for 10 min. The cell pellet is then frozen in liquid nitrogen. Pellets usually contain around  $10^9$  cells. For membrane isolation, pellets are freeze/thawed 3 times. They are then resuspended in ice cold 10mM Tris (pH 7.5), 1 mM EDTA (sodium salt) (40 mls/1e8

cells) and homogenized using a Dounce (glass/glass) homogenizer (20-25 strokes), followed by a Polytron suspension with 3--10 sec pulses on a  $\frac{3}{4}$  setting (Brinkman tissue homogenizer). This suspension is centrifuged at 300 x g for 10 min. The pellet is discarded and the supernatant fraction is centrifuged at 40,000 x g (Sorvall SS-34: 18,000 rpm) for 30 min. at 4 °C, resuspended in homogenizing buffer using the polytron, and washed one time. The pellet is resuspended in assay buffer (50 mM Tris pH 7.5) at a concentration of 1 - 4 mg protein/ml.

Membranes obtained this way are suitable for the set-up of a high throughput fatty acid ligand binding competition assay to search for compounds that interfere in the ligand-receptor interaction. The total binding of a labelled fatty acid to these membranes is first tested to be linear with the amount of membranes used. The time period to reach equilibrium binding at a suitable temperature is also established and is in our experience about 1 h at a temperature of 20 °C. Conditions for such a screening assay are exemplified by those used in MCH receptor binding. Typically 25 µg of membrane protein per well is used in a total volume of 100 µl buffer containing 50 mM HEPES, 1 mM CaCl<sub>2</sub>, 5 mM MgCl<sub>2</sub> and 0.5 % Bovine serum albumin (western blot quality), pH 7.4. The concentration of <sup>125</sup>I-mMCH is typically 1-2 nM and 75,000 cpm /well. Specific binding of <sup>125</sup>I-mMCH should be displaced completely by unlabeled MCH at concentrations of 100 nM or more.

The compounds to be tested are typically dissolved and added in DMSO and final concentrations of DMSO in the assay are 1 % or less. After incubation the contents of the wells are harvested on a polyethyleneimine-treated GF/C filter using a 96 well plate cell harvester and the filters are washed four times with typically 1 ml icecold wash buffer containing 20 mM HEPES 0.5M NaCl pH 7.4. The filters are counted to identify antagonists of labelled fatty acid ligand binding.

That which is claimed is:

1. A method of screening a test compound to determine whether the compound is a GPR40 receptor ligand, comprising detecting whether said test compound competitively inhibits the binding of a fatty acid GPR40 ligand to a GPR40 receptor.

2. A method according to claim 1 where said fatty acid is labeled with a detectable label.

3. A method according to claim 1 where said GPR40 receptor is provided by cell membranes containing functional GPR40 receptors.

4. A method according to claim 1 where said fatty acid GPR40 ligand is selected from Table 4 herein.

5. A method of screening a compound for GPR40 antagonist activity, comprising:  
(a) providing a test cell expressing on its surface a GPR40 receptor, and containing a reporter construct comprising a Gq responsive transcriptional element and a reporter gene, where expression of the reporter gene produces a detectable signal;

(b) exposing said test cell surface to a GPR40 fatty acid agonist under conditions that permit binding of a GPR40 fatty acid agonist;

(c) exposing said test cell surface to a test compound,

(d) measuring any detectable signal produced by said reporter gene, and

(e) comparing the detectable signal measured in (d) to that expected to occur in the presence of said GPR40 fatty acid agonist;

where a decrease in the detectable signal, compared to the detectable signal expected to occur due to the fatty acid agonist, indicates said test compound is a GPR40 antagonist.

6. a method according to claim 5 where said comparing step (e) comprises comparing the measured detectable signal to a pre-determined standard.

7. a method according to claim 5 where said comparing step (e) comprises providing a control cell, exposing said control cell to said GPR40 fatty acid agonist, measuring the detectable signal produced by the reporter gene, and comparing said control cell measurements with said test cell measurements.

8. A method according to claim 5 where said reporter gene encodes an enzyme selected from luciferase, phosphatase, secreted alkaline phosphatase,  $\beta$ -galactosidase, green fluorescent protein, beta lactamase, and chloramphenicol acetyl transferase.

9. A method according to claim 5 where said cell is a mammalian cell.

10. A method of screening a compound for GPR40 agonist activity, comprising:  
(a) providing a cell expressing on its surface a GPR40 receptor, and containing a reporter construct comprising:

- (i) a responsive transcriptional element selected from a Gi responsive transcriptional element, a Gq responsive transcriptional element, and a Gi/Gq responsive transcriptional element; and
- (ii) a reporter gene, where expression of the reporter gene produces a detectable signal;

(b) contacting a test compound to said cell surface, under conditions that permit binding of a GPR40 ligand to the GPR40 receptor, and

(c) detecting any reporter gene expression.

11. A method according to claim 10 where said reporter gene encodes an enzyme selected from luciferase, alkaline phosphatase, secreted alkaline phosphatase,  $\beta$ -galactosidase, green fluorescent protein, beta lactamase, and chloramphenicol acetyl transferase.

12. A method according to claim 10 where said cell is a mammalian cell.

13. A method according to claim 10 where said responsive transcriptional element is a Gi responsive transcriptional element.

14. A method according to claim 10 where said responsive transcriptional element is a Gq responsive transcriptional element.

15. A method according to claim 10 where said responsive transcriptional element is a Gi/Gq responsive transcriptional element.

16. A method according to claim 15 where said Gi/Gq responsive transcriptional element is Gal4/ELK-1.

17. A method of screening a compound for GPR40 antagonist activity, comprising:

(a) providing a cell expressing on its surface a GPR40 receptor, and containing a reporter construct comprising:

- (i) a responsive transcriptional element selected from a Gi responsive transcriptional element, a Gq responsive transcriptional element, and a Gi/Gq responsive transcriptional element; and
- (ii) a reporter gene, where expression of the reporter gene produces a detectable signal;

(b) exposing said cell surface to a test compound and a GPR40 agonist under conditions that permit binding of a GPR40 ligand to the GPR40 receptor; and

(c) detecting reporter gene expression;

where decreased reporter gene expression in the presence of both test compound and agonist, compared to reporter gene expression in the presence of said GPR40 agonist only, indicates that said test compound is a GPR40 antagonist.

18. A method according to claim 17 where said reporter gene encodes an enzyme selected from luciferase, phosphatase, secreted alkaline phosphatase,  $\beta$ -galactosidase,

green fluorescent protein, beta lactamase, and chloramphenicol acetyl transferase.

19. A method according to claim 17 where said cell is a mammalian cell.

20. A method according to claim 17 where said responsive transcriptional element is a Gi responsive transcriptional element.

21. A method according to claim 17 where said responsive transcriptional element is a Gq responsive transcriptional element.

22. A method according to claim 17 where said responsive transcriptional element is a Gi/Gq responsive transcriptional element.

23. A method according to claim 22 where said Gi/Gq responsive transcriptional element is Gal4/ELK-1.

24. A method of screening a compound for GPR40 antagonist activity, comprising detecting whether said compound decreases glucose-stimulated insulin release from mammalian pancreatic beta cells in the presence of a GPR40 agonist, compared to glucose-stimulated insulin release that would occur due to the presence of the GPR40 agonist.

25. A method according to claim 24 wherein said GPR40 agonist is a fatty acid.


26. A method according to claim 24 where said fatty acid is labeled with a detectable label.

27. A method of screening a compound for GPR40 agonist activity, comprising detecting whether said compound binds to GPR40 and increases glucose-stimulated insulin release from mammalian pancreatic beta cells.


28. A method according to claim 27 comprising:

- (a) providing a mammalian pancreatic cell expressing on its surface a GPR40 receptor,
- (b) contacting a test compound to said cell surface, under conditions that permit binding of GPR40 ligands to GPR40 receptors, and
- (c) detecting whether said test compound increases glucose-stimulated insulin release from said cell, compared to that which would be expected in the absence of a GPR40 agonist.

1/6

**Figure 1**

2/6



3/6

Fig. 3A





Fig 3B



4/6

Fig 3c




Fig. 4A




Fig. 4B





Fig. 4C



6/6

Fig. 4D





Figure 4E



1/8  
SEQUENCE LISTING

<110> SmithKline Beecham p.l.c.

Celia , Briscoe P

Diane , Ignar M

Alison, Muir I

Tadayyon, Mohammed

<120> Method of Screening for GPR40 Ligands

<130> tbd

<160> 10

<170> PatentIn version 3.1

<210> 1

<211> 903

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(900)

<223>

<400> 1

atg	gac	ctg	ccc	ccg	cag	ctc	tcc	ttc	ggc	ctc	tat	gtg	gcc	gcc	ttt	48
Met	Asp	Leu	Pro	Pro	Gln	Leu	Ser	Phe	Gly	Leu	Tyr	Val	Ala	Ala	Phe	
1				5					10				15			

gcg	ctg	ggc	ttc	ccg	ctc	aac	gtc	ctg	gcc	atc	cga	ggc	gcg	acg	gcc	96
Ala	Leu	Gly	Phe	Pro	Leu	Asn	Val	Leu	Ala	Ile	Arg	Gly	Ala	Thr	Ala	
		20					25					30				

cac	gcc	cgg	ctc	cgt	ctc	acc	cct	agc	ctg	gtc	tac	gcc	ctg	aac	ctg	144
His	Ala	Arg	Leu	Arg	Leu	Thr	Pro	Ser	Leu	Val	Tyr	Ala	Leu	Asn	Leu	
		35				40						45				

ggc	tgc	tcc	gac	ctg	ctg	ctg	aca	gtc	tct	ctg	ccc	ctg	aag	gcg	gtg	192
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

Gly	Cys	Ser	Asp	Leu	Leu	Leu	Thr	Val	Ser	Leu	Pro	Leu	Lys	Ala	Val	
50						55					60					
gag	gcg	cta	gcc	tcc	ggg	gcc	tgg	cct	ctg	ccg	gcc	tcg	ctg	tgc	ccc	240
Glu	Ala	Leu	Ala	Ser	Gly	Ala	Trp	Pro	Leu	Pro	Ala	Ser	Leu	Cys	Pro	
65					70					75					80	
gtc	ttc	gcg	gtg	gcc	cac	ttc	ttc	cca	ctc	tat	gcc	ggc	ggg	ggc	ttc	288
Val	Phe	Ala	Val	Ala	His	Phe	Phe	Pro	Leu	Tyr	Ala	Gly	Gly	Gly	Phe	
				85					90					95		
ctg	gcc	gcc	ctg	agt	gca	ggc	cgc	tac	ctg	gga	gca	gcc	ttc	ccc	ttg	336
Leu	Ala	Ala	Leu	Ser	Ala	Gly	Arg	Tyr	Leu	Gly	Ala	Ala	Phe	Pro	Leu	
			100					105					110			
ggc	tac	caa	gcc	ttc	cgg	agg	ccg	tgc	tat	tcc	tgg	ggg	gtg	tgc	gcg	384
Gly	Tyr	Gln	Ala	Phe	Arg	Arg	Pro	Cys	Tyr	Ser	Trp	Gly	Val	Cys	Ala	
		115					120					125				
gcc	atc	tgg	gcc	ctc	gtc	ctg	tgt	cac	ctg	ggg	ctg	gtc	ttt	ggg	ttg	432
Ala	Ile	Trp	Ala	Leu	Val	Leu	Cys	His	Leu	Gly	Leu	Val	Phe	Gly	Leu	
	130					135					140					
gag	gct	cca	gga	ggc	tgg	ctg	gac	cac	agc	aac	acc	tcc	ctg	ggc	atc	480
Glu	Ala	Pro	Gly	Gly	Trp	Leu	Asp	His	Ser	Asn	Thr	Ser	Leu	Gly	Ile	
145					150					155					160	
aac	aca	ccg	gtc	aac	ggc	tct	ccg	gtc	tgc	ctg	gag	gcc	tgg	gac	ccg	528
Asn	Thr	Pro	Val	Asn	Gly	Ser	Pro	Val	Cys	Leu	Glu	Ala	Trp	Asp	Pro	
				165					170					175		
gcc	tct	gcc	ggc	ccg	gcc	cgc	ttc	agc	ctc	tct	ctc	ctg	ctc	ttt	ttt	576
Ala	Ser	Ala	Gly	Pro	Ala	Arg	Phe	Ser	Leu	Ser	Leu	Leu	Leu	Phe	Phe	
			180					185					190			
ctg	ccc	ttg	gcc	atc	aca	gcc	ttc	tgc	tac	gtg	ggc	tgc	ctc	cgg	gca	624
Leu	Pro	Leu	Ala	Ile	Thr	Ala	Phe	Cys	Tyr	Val	Gly	Cys	Leu	Arg	Ala	
		195					200					205				
ctg	gcc	cgc	tcc	ggc	ctg	acg	cac	agg	cgg	aag	ctg	cgg	gcc	gcc	tgg	672
Leu	Ala	Arg	Ser	Gly	Leu	Thr	His	Arg	Arg	Lys	Leu	Arg	Ala	Ala	Trp	
	210					215					220					
gtg	gcc	ggc	ggg	gcc	ctc	ctc	acg	ctg	ctg	ctc	tgc	gta	gga	ccc	tac	720
Val	Ala	Gly	Gly	Ala	Leu	Leu	Thr	Leu	Leu	Leu	Cys	Val	Gly	Pro	Tyr	
225					230					235					240	
aac	gcc	tcc	aac	gtg	gcc	agc	ttc	ctg	tac	ccc	aat	cta	gga	ggc	tcc	768
Asn	Ala	Ser	Asn	Val	Ala	Ser	Phe	Leu	Tyr	Pro	Asn	Leu	Gly	Gly	Ser	
				245					250					255		
tgg	cgg	aag	ctg	ggg	ctc	atc	acg	ggg	gcc	tgg	agt	gtg	gtg	ctt	aat	816
Trp	Arg	Lys	Leu	Gly	Leu	Ile	Thr	Gly	Ala	Trp	Ser	Val	Val	Leu	Asn	
			260					265					270			
ccg	ctg	gtg	acc	ggg	tac	ttg	gga	agg	ggg	cct	ggc	ctg	aag	aca	gtg	864
Pro	Leu	Val	Thr	Gly	Tyr	Leu	Gly	Arg	Gly	Pro	Gly	Leu	Lys	Thr	Val	
		275					280					285				

3/8

tgt gcg gca aga acg caa ggg ggc aag tcc cag aag taa  
 Cys Ala Ala Arg Thr Gln Gly Gly Lys Ser Gln Lys  
       290                      295                      300

903

&lt;210&gt; 2

&lt;211&gt; 300

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 2

Met Asp Leu Pro Pro Gln Leu Ser Phe Gly Leu Tyr Val Ala Ala Phe  
 1                      5                      10                      15  
 Ala Leu Gly Phe Pro Leu Asn Val Leu Ala Ile Arg Gly Ala Thr Ala  
                       20                      25                      30  
 His Ala Arg Leu Arg Leu Thr Pro Ser Leu Val Tyr Ala Leu Asn Leu  
                       35                      40                      45  
 Gly Cys Ser Asp Leu Leu Leu Thr Val Ser Leu Pro Leu Lys Ala Val  
                       50                      55                      60  
 Glu Ala Leu Ala Ser Gly Ala Trp Pro Leu Pro Ala Ser Leu Cys Pro  
 65                      70                      75                      80  
 Val Phe Ala Val Ala His Phe Phe Pro Leu Tyr Ala Gly Gly Gly Phe  
                       85                      90                      95  
 Leu Ala Ala Leu Ser Ala Gly Arg Tyr Leu Gly Ala Ala Phe Pro Leu  
                       100                      105                      110  
 Gly Tyr Gln Ala Phe Arg Arg Pro Cys Tyr Ser Trp Gly Val Cys Ala  
                       115                      120                      125  
 Ala Ile Trp Ala Leu Val Leu Cys His Leu Gly Leu Val Phe Gly Leu  
                       130                      135                      140  
 Glu Ala Pro Gly Gly Trp Leu Asp His Ser Asn Thr Ser Leu Gly Ile  
 145                      150                      155                      160  
 Asn Thr Pro Val Asn Gly Ser Pro Val Cys Leu Glu Ala Trp Asp Pro  
                       165                      170                      175  
 Ala Ser Ala Gly Pro Ala Arg Phe Ser Leu Ser Leu Leu Leu Phe Phe  
                       180                      185                      190  
 Leu Pro Leu Ala Ile Thr Ala Phe Cys Tyr Val Gly Cys Leu Arg Ala  
                       195                      200                      205  
 Leu Ala Arg Ser Gly Leu Thr His Arg Arg Lys Leu Arg Ala Ala Trp  
                       210                      215                      220  
 Val Ala Gly Gly Ala Leu Leu Thr Leu Leu Leu Cys Val Gly Pro Tyr  
 225                      230                      235                      240

4/8

Asn	Ala	Ser	Asn	Val	Ala	Ser	Phe	Leu	Tyr	Pro	Asn	Leu	Gly	Gly	Ser
				245					250					255	
Trp	Arg	Lys	Leu	Gly	Leu	Ile	Thr	Gly	Ala	Trp	Ser	Val	Val	Leu	Asn
			260					265					270		
Pro	Leu	Val	Thr	Gly	Tyr	Leu	Gly	Arg	Gly	Pro	Gly	Leu	Lys	Thr	Val
		275					280					285			
Cys	Ala	Ala	Arg	Thr	Gln	Gly	Gly	Lys	Ser	Gln	Lys				
	290					295					300				

<210> 3

<211> 903

<212> DNA

<213> Mus musculus

<220>

<221> CDS

<222> (1) . . (900)

<223>

<400> 3

atg gac ctg ccc cca cag ctc tcc ttc gct ctc tat gta tct gcc ttt 48  
Met Asp Leu Pro Pro Gln Leu Ser Phe Ala Leu Tyr Val Ser Ala Phe  
1 5 10 15

gcg ctg ggc ttt cca ttg aac ttg tta gcc atc cga ggc gca gtg tcc 96  
Ala Leu Gly Phe Pro Leu Asn Leu Leu Ala Ile Arg Gly Ala Val Ser  
20 25 30

cac gct aaa ctg cga ctc act ccc agc ttg gtc tac act ctc cat ctg      144  
His Ala Lys Leu Arg Leu Thr Pro Ser Leu Val Tyr Thr Leu His Leu  
          35                       40                       45

ggc tgc tct gat ctc cta ctg gcc atc act ctg ccc ctg aag gct gtg 192  
Gly Cys Ser Asp Leu Leu Leu Ala Ile Thr Leu Pro Leu Lys Ala Val  
50 55 60

gag gcc ctg gct tct gga gcc tgg ccc ctg ccg ctc ccc ttc tgc cca 240  
Glu Ala Leu Ala Ser Gly Ala Trp Pro Leu Pro Leu Pro Phe Cys Pro  
65 70 75 80

gtc ttt gcc ttg gcc cac ttt gct ccc ctc tac gca ggc gga ggc ttc 288  
Val Phe Ala Leu Ala His Phe Ala Pro Leu Tyr Ala Gly Gly Gly Phe  
85 90 95

cta gct gct ctc agc gct ggc cgc tac ctg ggg gct gcc ttc ccc ttc 336  
Leu Ala Ala Leu Ser Ala Gly Arg Tyr Leu Gly Ala Ala Phe Pro Phe  
100 105 110

```

ggg tac caa gcc atc cgg agg ccc cgc tat tcc tgg ggt gtg tgt gtg      384
Gly Tyr Gln Ala Ile Arg Arg Pro Arg Tyr Ser Trp Gly Val Cys Val
      115                      120                      125

gct ata tgg gcc ctt gtc ctc tgc cac ctg ggg ctg gcc ctt ggc ttg      432
Ala Ile Trp Ala Leu Val Leu Cys His Leu Gly Leu Ala Leu Gly Leu
      130                      135                      140

gag act tcc gga agc tgg ctg gac aac agt acc agt tcc ctg ggc atc      480
Glu Thr Ser Gly Ser Trp Leu Asp Asn Ser Thr Ser Ser Leu Gly Ile
      145                      150                      155                      160

aac ata ccc gtg aat ggc tcc ccg gtc tgc ctg gaa gcc tgg gat ccc      528
Asn Ile Pro Val Asn Gly Ser Pro Val Cys Leu Glu Ala Trp Asp Pro
      165                      170                      175

gac tct gcc cgc cct gcc cgt ctc agt ttc tcc att ctg ctc ttc ttt      576
Asp Ser Ala Arg Pro Ala Arg Leu Ser Phe Ser Ile Leu Leu Phe Phe
      180                      185                      190

ctg ccc ttg gtc atc act gcc ttc tgc tat gtg ggc tgc ctc cgg gcc      624
Leu Pro Leu Val Ile Thr Ala Phe Cys Tyr Val Gly Cys Leu Arg Ala
      195                      200                      205

ctg gtg cgc tca ggc ctg agc cac aaa cgg aag ctc agg gca gct tgg      672
Leu Val Arg Ser Gly Leu Ser His Lys Arg Lys Leu Arg Ala Ala Trp
      210                      215                      220

gtg gcc gga ggc gct ctc ctc aca ctc ctg ctc tgc ctg ggg ccc tat      720
Val Ala Gly Gly Ala Leu Leu Thr Leu Leu Leu Cys Leu Gly Pro Tyr
      225                      230                      235                      240

aat gcc tcc aat gtg gct agt ttc ata aac ccg gac cta gga ggc tcc      768
Asn Ala Ser Asn Val Ala Ser Phe Ile Asn Pro Asp Leu Gly Gly Ser
      245                      250                      255

tgg agg aag ttg gga ctc atc aca ggg gcc tgg agt gtg gta ctc aac      816
Trp Arg Lys Leu Gly Leu Ile Thr Gly Ala Trp Ser Val Val Leu Asn
      260                      265                      270

cca ctg gtc act ggc tac ttg gga aca ggt cct gga cgg gga aca ata      864
Pro Leu Val Thr Gly Tyr Leu Gly Thr Gly Pro Gly Arg Gly Thr Ile
      275                      280                      285

tgt gtg acg agg act caa aga gga aca att cag aag tag      903
Cys Val Thr Arg Thr Gln Arg Gly Thr Ile Gln Lys
      290                      295                      300

```

&lt;210&gt; 4

&lt;211&gt; 300

&lt;212&gt; PRT

&lt;213&gt; Mus musculus

&lt;400&gt; 4

Met	Asp	Leu	Pro	Pro	Gln	Leu	Ser	Phe	Ala	Leu	Tyr	Val	Ser	Ala	Phe	1	5	10	15
Ala	Leu	Gly	Phe	Pro	Leu	Asn	Leu	Leu	Ala	Ile	Arg	Gly	Ala	Val	Ser	20	25	30	
His	Ala	Lys	Leu	Arg	Leu	Thr	Pro	Ser	Leu	Val	Tyr	Thr	Leu	His	Leu	35	40	45	
Gly	Cys	Ser	Asp	Leu	Leu	Leu	Ala	Ile	Thr	Leu	Pro	Leu	Lys	Ala	Val	50	55	60	
Glu	Ala	Leu	Ala	Ser	Gly	Ala	Trp	Pro	Leu	Pro	Leu	Pro	Phe	Cys	Pro	65	70	75	80
Val	Phe	Ala	Leu	Ala	His	Phe	Ala	Pro	Leu	Tyr	Ala	Gly	Gly	Gly	Phe	85	90	95	
Leu	Ala	Ala	Leu	Ser	Ala	Gly	Arg	Tyr	Leu	Gly	Ala	Ala	Phe	Pro	Phe	100	105	110	
Gly	Tyr	Gln	Ala	Ile	Arg	Arg	Pro	Arg	Tyr	Ser	Trp	Gly	Val	Cys	Val	115	120	125	
Ala	Ile	Trp	Ala	Leu	Val	Leu	Cys	His	Leu	Gly	Leu	Ala	Leu	Gly	Leu	130	135	140	
Glu	Thr	Ser	Gly	Ser	Trp	Leu	Asp	Asn	Ser	Thr	Ser	Ser	Leu	Gly	Ile	145	150	155	160
Asn	Ile	Pro	Val	Asn	Gly	Ser	Pro	Val	Cys	Leu	Glu	Ala	Trp	Asp	Pro	165	170	175	
Asp	Ser	Ala	Arg	Pro	Ala	Arg	Leu	Ser	Phe	Ser	Ile	Leu	Leu	Phe	Phe	180	185	190	
Leu	Pro	Leu	Val	Ile	Thr	Ala	Phe	Cys	Tyr	Val	Gly	Cys	Leu	Arg	Ala	195	200	205	
Leu	Val	Arg	Ser	Gly	Leu	Ser	His	Lys	Arg	Lys	Leu	Arg	Ala	Ala	Trp	210	215	220	
Val	Ala	Gly	Gly	Ala	Leu	Leu	Thr	Leu	Leu	Leu	Cys	Leu	Gly	Pro	Tyr	225	230	235	240
Asn	Ala	Ser	Asn	Val	Ala	Ser	Phe	Ile	Asn	Pro	Asp	Leu	Gly	Gly	Ser	245	250	255	
Trp	Arg	Lys	Leu	Gly	Leu	Ile	Thr	Gly	Ala	Trp	Ser	Val	Val	Leu	Asn	260	265	270	
Pro	Leu	Val	Thr	Gly	Tyr	Leu	Gly	Thr	Gly	Pro	Gly	Arg	Gly	Thr	Ile	275	280	285	
Cys	Val	Thr	Arg	Thr	Gln	Arg	Gly	Thr	Ile	Gln	Lys	290	295	300					

<210> 5

<211> 20

<212> DNA

<213> Forward primer for human GPR40

<400> 5

gtggtgctta atccgctggt

20

<210> 6

<211> 21

<212> DNA

<213> Reverse primer for human GPR40

<400> 6

tggcggttact tctgggactt g

21

<210> 7

<211> 25

<212> DNA

<213> Probe for human GPR40

<400> 7

cttgcgttct tgccgcacac actgt

25

<210> 8

<211> 21

<212> DNA

<213> Forward primer for mouse/rat GPR40

<400> 8

agttccctgg gcatcaacat a

21

<210> 9

<211> 22

<212> DNA

<213> Reverse primer for mouse/rat GPR40

<400> 9

caagggcaga aagaagagca ga

22

<210> 10

<211> 22

<212> DNA

<213> Probe for mouse/rat GPR40

<400> 10

aatggctccc cggtctgcct gg

22

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
25 July 2002 (25.07.2002)

PCT

(10) International Publication Number  
**WO 02/057783 A3**

(51) International Patent Classification<sup>7</sup>: **G01N 33/566**,  
C12Q 1/68, G01N 33/50, 33/74

Research Triangle Park, NC 27709 (US). **TADAYYON, Mohammad** [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB).

(21) International Application Number: PCT/US01/48985

(74) Agents: **LEVY, David, J.** et al.; GlaxoSmithKline, Five Moore Drive, PO Box 13398, Research Triangle Park, NC 27709 (US).

(22) International Filing Date:  
18 December 2001 (18.12.2001)

(25) Filing Language: English

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(26) Publication Language: English

(30) Priority Data:  
0031527.5 22 December 2000 (22.12.2000) GB

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicants (*for all designated States except US*): **GLAXO GROUP LIMITED** [GB/GB]; Glaxo Wellcome House, Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB). **SMITHKLINE BEECHAM P.L.C.** [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **ANDREWS, John, Leon** [US/US]; GlaxoSmithKline, Five Moore Drive, PO Box 13398, Research Triangle Park, NC 27709 (US). **BRISCOE, Celia, Patricia** [GB/US]; GlaxoSmithKline, Five Moore Drive, PO Box 13398, Research Triangle Park, NC 27709 (US). **IGNAR, Diane, Michelle** [US/US]; GlaxoSmithKline, Five Moore Drive, PO Box 13398, Research Triangle Park, NC 27709 (US). **MUIR, Allison, Isobel** [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB). **SAULS, Howard, Ray Jr.** [US/US]; GlaxoSmithKline, Five Moore Drive, PO Box 13398,

Published:

— with international search report

(88) Date of publication of the international search report:  
7 August 2003

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: METHOD OF SCREENING FOR GPR40 LIGANDS

(57) Abstract: Methods of screening for ligands of the GPR40 receptor are provided, including methods utilizing fatty acid GPR40 ligands.



**WO 02/057783 A3**

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/48985

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 G01N33/566 C12Q1/68 G01N33/50 G01N33/74

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, PAJ, WPI Data, BIOSIS, EMBASE, SEQUENCE SEARCH, EMBL

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 00 22129 A (ARENA PHARMACEUTICALS INC ;LIAW CHEN W (US); BEHAN DOMINIC P (US);) 20 April 2000 (2000-04-20) page 24, line 20 -page 26, line 15; table A page 37, line 8 - line 20 page 65, line 11 -page 66, line 13; claim 32</p> <p style="text-align: center;">--- -/--</p>	10-24, 27, 28



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*Z\* document member of the same patent family

Date of the actual completion of the international search

18 March 2003

Date of mailing of the international search report

23/04/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Vadot-Van Geldre, E

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 01/48985

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>SAWZDARGO M ET AL: "A CLUSTER OF FOUR NOVEL HUMAN G PROTEIN-COUPLED RECEPTOR GENES OCCURRING IN CLOSE PROXIMITY TO CD22 GENE ON CHROMOSOME 19Q13.1" BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, ACADEMIC PRESS INC. ORLANDO, FL, US, vol. 239, 1997, pages 543-547, XP002067401 ISSN: 0006-291X cited in the application page 543</p> <p>----</p>	10-23
Y	<p>STABLES J ET AL: "DEVELOPMENT OF A DUAL GLOW-SIGNAL FIREFLY AND RENILLA LUCIFERASE ASSAY REAGENT FOR THE ANALYSIS OF G-PROTEIN COUPLED RECEPTOR SIGNALLING" JOURNAL OF RECEPTOR AND SIGNAL TRANSDUCTION RESEARCH, MARCEL DEKKER, NEW YORK, NY, US, vol. 19, no. 1-4, 1999, pages 395-410, XP000982483 ISSN: 1079-9893 cited in the application abstract page 395</p> <p>----</p>	10-23
Y	<p>SHARIF T R ET AL: "A HIGH THROUGHPUT SYSTEM FOR THE EVALUATION OF PROTEIN KINASE C INHIBITORS BASED ON ELK1 TRANSCRIPTIONAL ACTIVATION IN HUMAN ASTROCYTOMA CELLS" INTERNATIONAL JOURNAL OF ONCOLOGY, EDITORIAL ACADEMY OF THE INTERNATIONAL JOURNAL OF ONCOLOGY,, GR, vol. 14, no. 2, February 1999 (1999-02), pages 326-335, XP001078793 ISSN: 1019-6439 abstract</p> <p>----</p>	10-23
X	<p>DATABASE EMBL 'Online! 20 March 2000 (2000-03-20) Database accession no. AW583167 XP002233764 abstract</p> <p>----</p>	24,27,28
E	<p>DATABASE WPI Section Ch, Week 200249 Derwent Publications Ltd., London, GB; Class B04, AN 2002-463740 XP002233765 &amp; WO 02 44362 A (YAMANOUCHI PHARM CO LTD), 6 June 2002 (2002-06-06) abstract</p> <p>----</p> <p style="text-align: center;">-/--</p>	1-28

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 01/48985

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WARNOTTE CATHERINE ET AL: "Unbound rather than total concentration and saturation rather than unsaturation determine the potency of fatty acids on insulin secretion." MOLECULAR AND CELLULAR ENDOCRINOLOGY, vol. 153, no. 1-2, 20 July 1999 (1999-07-20), pages 147-153, XP002233763 ISSN: 0303-7207 cited in the application abstract ---	1-28
A	TSENG C-C ET AL: "ROLE OF REGULATOR OF G PROTEIN SIGNALING IN DESENSITIZATION OF THE GLUCOSE-DEPENDENT INSULINOTROPIC PEPTIDE RECEPTOR" ENDOCRINOLOGY, BALTIMORE, MD, US, vol. 139, no. 11, November 1998 (1998-11), pages 4470-4475, XP000853560 ISSN: 0013-7227 page 4470 ---	1-28
A	EP 0 849 361 A (SMITHKLINE BEECHAM PLC ;SMITHKLINE BEECHAM CORP (US)) 24 June 1998 (1998-06-24) claims 11,12 -----	1-28

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/48985

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0022129	A	20-04-2000	AU 6299199 A	01-05-2000
			AU 6430799 A	01-05-2000
			CA 2342314 A1	20-04-2000
			CN 1398298 T	19-02-2003
			DE 1121431 T1	23-05-2002
			EP 1137776 A2	04-10-2001
			EP 1121431 A1	08-08-2001
			ES 2163384 T1	01-02-2002
			JP 2002527727 T	27-08-2002
			US 2003023069 A1	30-01-2003
			WO 0021987 A2	20-04-2000
			WO 0022129 A1	20-04-2000
			WO 0022131 A2	20-04-2000
			US 2003018182 A1	23-01-2003
			US 2002193584 A1	19-12-2002
<hr/>				
WO 0244362	A	06-06-2002	AU 1850902 A	11-06-2002
			WO 0244362 A1	06-06-2002
<hr/>				
EP 0849361	A	24-06-1998	US 6309854 B1	30-10-2001
			CA 2218452 A1	07-06-1998
			DE 849361 T1	25-01-2001
			EP 0849361 A2	24-06-1998
			JP 10229887 A	02-09-1998
			US 2002082202 A1	27-06-2002
			US 6001963 A	14-12-1999
<hr/>				